



# AQUATHOL® K

## AQUATIC HERBICIDE

**ACTIVE INGREDIENT:**

Dipotassium salt of endosulfar\* ..... 40.3%

OTHER INGREDIENTS: ..... 59.7%

TOTAL ..... 100.0%

\*7-oxabicyclo [2.2.1]heptane-2,3-dicarboxylic acid equivalent 28.6%

Contains per gallon 4.23 lbs. dipotassium endosulfar

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

**FIRST AID:**

**IF IN EYES:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

**IF INHALED:**

- Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow. Do not induce vomiting unless told by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (303) 623-5716 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** Measures against circulatory shock, respiratory depression, and convulsion may be needed.

EPA Registration No. 4581-204-82695    EPA Establishment No. 62171-MS-003

Net Contents \_\_\_\_\_

Sold by:  
Cerexagri-Nisso LLC  
630 Freedom Business Center • Suite 402  
King of Prussia, PA 19406  
1 800-438-6071 • www.cerexagri-nisso.com



Cerexagri-Nisso LLC

## **GENERAL INFORMATION**

AQUATHOL K is a liquid concentrate soluble in water which is effective against a broad range of aquatic plants with a margin of safety to fish.

Dosage rates indicated for the application of AQUATHOL K are measured in "Parts Per Million" (ppm) of dipotassium endothall. Only 0.5 to 5.0 ppm are generally required for aquatic weed control, whereas some fish species are tolerant to approximately 100 ppm or over.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### **HOW TO APPLY:**

AQUATHOL K is a contact herbicide; consequently, do not apply before weeds are present. Application as early as possible after weeds appear and are actively growing is recommended for best results.

If an entire pond is treated at one time, or if the dissolved oxygen level is low at time of application, decay of weeds may remove enough oxygen from the water, causing fish to suffocate. Water containing very heavy vegetation should be treated in sections to prevent suffocation of fish. Sections should be treated 5-7 days apart. Carefully measure size and depth of area to be treated and determine amount of AQUATHOL K to apply from chart.

AQUATHOL K should be sprayed on the water or injected below the water surface and should be distributed as evenly as possible. It may be applied as a concentrate or diluted with water depending on the equipment. Some dilution will give better distribution. For best results apply when water is quiescent and/or flows are minimal.

In instances where the weed(s) to be controlled is an exposed surface problem (i.e., some of the broad-leaved pond weeds) coverage is important. For best results apply the concentrate or with the least amount of water compatible with the application equipment.

Necessary approval and/or permits should be obtained in states where required.

## **AQUATIC WEEDS CONTROLLED AND DOSAGE RATE CHARTS**

AQUATHOL K is recommended for the control of the following aquatic weeds in irrigation and drainage canals, ponds and lakes at the rates indicated. Since the active ingredient is water soluble and tends to diffuse from the treated area, select the dosage rate applicable to the area to be treated. Use the lower rate in each range of rates where the growth is young and growing and/or where the weed stand is not heavy. Marginal treatments of large bodies of water require higher rates as indicated.

Aquatic Weed	RATES			
	Entire Pond/Lake or Large Area Treatment	Gallons per Acre Ft.	Spot or Lake Margin Treatment	Gallons per Acre Ft.
Bur Reed, Sparganium spp.	3.0-4.0 ppm	1.9-2.6 gal.	4.0-5.0 ppm	2.6-3.2 gal.
Coontail, Ceratophyllum spp.	1.0-2.0 ppm	0.6-1.3 gal.	2.0-3.0 ppm	1.3-1.9 gal.
Horned Pondweed, Zannichellia palustris	1.0-2.0 ppm	0.6-1.3 gal.	2.0-3.0 ppm	1.3-1.9 gal.
Hydrilla, Hydrilla verticillata	2.0-3.0 ppm	1.3-1.9 gal.	3.0-4.0 ppm	1.9-2.6 gal.
Hygrophila, Hygrophila polysperma	4.0-5.0 ppm	2.6-3.2 gal.	5.0 ppm	3.2 gal.
Milfoil, Myriophyllum spp.	2.0-3.0 ppm	1.3-1.9 gal.	3.0-4.0 ppm	1.9-2.6 gal.
Naiad, Najas spp.	1.0-3.0 ppm	0.6-1.9 gal.	2.0-4.0 ppm	1.3-2.6 gal.
Pondweed, Potamogeton spp.	0.5-3.0 ppm	0.3-1.9 gal.	1.5-4.0 ppm	1.0-2.6 gal.
Including:				
American, P. nodosus	2.0-3.0 ppm	1.3-1.9 gal.	3.0-4.0 ppm	1.9-2.6 gal.
Largeleaf (Bass Weed), P. amplifolius	2.0-3.0 ppm	1.3-1.9 gal.	3.0-4.0 ppm	1.9-2.6 gal.
Curlyleaf, P. crispus	0.5-1.5 ppm	0.3-1.0 gal.	1.5-3.0 ppm	1.0-1.9 gal.
Flatstem, P. zosteriformis	2.0-3.0 ppm	1.3-1.9 gal.	3.0-4.0 ppm	1.9-2.6 gal.
Floating-leaf, P. natans	1.0-2.0 ppm	0.6-1.3 gal.	2.0-3.0 ppm	1.3-1.9 gal.
Illinois, P. illinoensis	1.5-2.5 ppm	1.0-1.6 gal.	2.5-3.5 ppm	1.6-2.3 gal.
Narrowleaf, P. pusillus	1.0-2.0 ppm	0.6-1.3 gal.	2.0-3.0 ppm	1.3-1.9 gal.
Threadleaf, P. filiformis	2.0-3.0 ppm	1.3-1.9 gal.	3.0-4.0 ppm	1.9-2.6 gal.
Sago, P. pectinatus	1.0-2.0 ppm	0.6-1.3 gal.	2.0-3.0 ppm	1.3-1.9 gal.
Variable Leaf, P. diversifolius	1.0-2.0 ppm	0.6-1.3 gal.	2.0-3.0 ppm	1.3-1.9 gal.
Parrot Feather, Myriophyllum aquaticum	2.0-3.0 ppm	1.3-1.9 gal.	3.0-4.0 ppm	1.9-2.6 gal.
Water Stargrass, Heteranthera spp.	2.0-3.0 ppm	1.3-1.9 gal.	3.0-4.0 ppm	1.9-2.6 gal.

**RATE OF APPLICATION — LAKES AND PONDS**

The following chart indicates the total quantity of material to be applied.

**APPROXIMATE GALLONS OF AQUATHOL K FOR ONE ACRE (208' x 208') TREATMENT**

DEPTH . . . . .	DOSAGE IN GALLONS FOR VARIOUS CONCENTRATIONS IN PPM						
	0.5 ppm	1.0 ppm	1.5 ppm	2.0 ppm	3.0 ppm	4.0 ppm	5.0 ppm
1 ft. . . . .	0.3	0.6	1.0	1.3	1.9	2.6	3.2
2 ft. . . . .	0.6	1.3	1.9	2.6	3.8	5.1	6.4
4 ft. . . . .	1.3	2.6	3.8	5.1	7.7	10.2	12.8
6 ft. . . . .	1.9	3.8	5.8	7.6	11.5	15.3	19.2

**RATE OF APPLICATION — IRRIGATION AND DRAINAGE CANALS\*\***

The following indicates the total quantity of material to be applied.

**GALLONS OF AQUATHOL K REQUIRED TO TREAT 1 MILE OF CANAL 1 FOOT DEEP\***

PPM	WIDTH OF CANAL IN FEET			
	5	10	15	20
1.0 ppm	0.4	0.75	1.2	1.5
2.0 ppm	0.75	1.5	2.3	3.0
3.0 ppm	1.2	2.3	3.5	4.5
4.0 ppm	1.5	3.0	4.5	6.0
5.0 ppm	2.0	3.8	5.7	7.5

The minimum contact time with weeds for optimum results should be 2 hours.

\*For deeper water, adjust rate accordingly.

\*\* Not for this use in California.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)**

**DANGER**

CORROSIVE. CAUSES IRREVERSIBLE EYE DAMAGE. MAY BE FATAL IF SWALLOWED. HARMFUL IF INHALED OR ABSORBED THROUGH SKIN. DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. AVOID BREATHING VAPORS OR SPRAY MIST. Wear appropriate protective eyewear such as goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- socks
- shoes
- chemical resistant gloves
- protective eyewear

**USER SAFETY RECOMMENDATIONS:**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove protective clothing and equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Avoid contact with or drift to other crops or plants as injury may result. Wash out spray equipment with water after each operation.

Not for use in brackish or salt water. Do not use water from treated areas for watering livestock, for preparing agricultural sprays for food crops, for irrigation or for domestic purposes within the following periods:

Up to 0.5 ppm dipotassium salt — 7 days after application

Up to 4.25 ppm dipotassium salt — 14 days after application

Up to 5.0 ppm dipotassium salt — 25 days after application

Treated water can be used for sprinkling bent grass immediately.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage Instructions:** Store in the original container. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. Storage at temperatures below 32°F may result in the product freezing or crystallizing. Should this occur the product must be warmed to 50°F or higher and thoroughly agitated. In the event of a spillage during handling or storage, absorb with sand or other inert material and dispose of absorbent in accordance with the Pesticide Disposal Instructions listed below.

**Pesticide Disposal Instructions:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal Instructions:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**EMERGENCY TELEPHONE NUMBERS:**

**CHEMTREC: (800) 424-9300**

**MEDICAL: (303) 623-5716 Rocky Mountain Poison Control Center**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, resistant strains or other influencing factors in the use of the product. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Cerexagri-Nisso LLC, Manufacturer and Seller harmless for any claims relating to such factors. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CEREXAGRI-NISSO LLC AND MANUFACTURER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THE LABEL.

To the extent consistent with applicable law, Cerexagri-Nisso LLC, Manufacturer or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CEREXAGRI-NISSO LLC, MANUFACTURER AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CEREXAGRI-NISSO LLC, MANUFACTURER OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Cerexagri-Nisso LLC, Manufacturer and Seller offer this product, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of Cerexagri-Nisso LLC.

*Aquathol is a registered trademark of Cerexagri, Inc.*

© 2006 Cerexagri, Inc. All rights reserved.

4581-204-82695(101906-1923)

Made and Printed in U.S.A.