

# Specimen Label



## Aquatic Herbicide

\*Trademark of SePRO Corporation

For control of floating, emersed, and submersed vegetation in still or flowing aquatic sites such as potable water sources, lakes, rivers, reservoirs, and ponds, slow-flowing or quiescent water bodies, crop and non-crop irrigation systems (canals, laterals, and ditches), fish, golf course, ornamental, swimming, and fire ponds and aquaculture including fish and shrimp.

### Active Ingredient:

Copper Carbonate*	15.9%
Inert Ingredients	84.1%
Total	100.0%

\*Metallic copper equivalent, 9.1%

### Keep Out of Reach of Children

## DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

**Note to Physician:** Probable mucosal damage may contraindicate the use of gastric lavage.

Refer to label booklet for additional precautionary information and Directions for Use, including Storage and Disposal.

**Notice:** Read the entire label. Use only according to label directions. **Before buying or using this product, read "Warranty Disclaimer", "Inherent Risks of Use" and "Limitation of Remedies" inside label booklet.**

For additional information on our products, please visit [www.sepro.com](http://www.sepro.com)

EPA Reg. No. 67690-10  
FPL 021005

EPA Est. No. 5905-GA-01  
SC-84-0042

\*Trademark of SePRO Corporation

SePRO Corporation • Carmel, IN 46032 U.S.A.

First Aid	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or a doctor for further treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call INFOTRAC 1-800-535-5053.</p>	

## Precautionary Statements

### Hazards to Humans and Domestic Animals

**DANGER:** Corrosive. Causes irreversible eye damage and skin burn. May be fatal if absorbed through skin. Harmful if swallowed. Do not get in eyes on skin or on clothing. Wear goggles, face shield, or safety glasses, protective clothing and chemical resistant gloves. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking and using tobacco. Remove contaminated clothing and wash before reuse.

Nautique\* Aquatic Herbicide

## Environmental Hazards

Fish toxicity is dependent on the hardness of the water. In soft water, trout and other species of fish may be killed at application rates recommended on this label. Do not use in waters containing trout or other sensitive species if the carbonate hardness of the water is less than 50 ppm. Fish toxicity generally decreases when the hardness of water increases. Do not treat more than one-half of lake or pond at one time to avoid depletion of oxygen levels due to decaying vegetation. Consult State Fish and Game Agency or other responsible Agency before applying this product to public waters.

---

## Directions for Use

---

It is a violation of Federal Law to use this product in a manner inconsistent with its label directions.

### General Information

Nautique may be applied to potable water sources, lakes, rivers, reservoirs, ponds, slow-flowing or quiescent water bodies, crop and non-crop irrigation systems (ditches, canals, and laterals), fish, golf course, ornamental, swimming, and fire ponds, and aquaculture including fish and shrimp. In waters with greater calcium carbonate hardness, the higher use rates are recommended for improved plant control.

### Target Species

Nautique Aquatic Herbicide is a double chelated copper formulation that provides effective control of floating, submersed, and emersed aquatic plants having a sensitivity to copper absorption including:

Coontail	Naiads
Curlyleaf Pondweed	Thin Leaf Pondweed
Egeria (Brazilian Elodia)	Vallisneria
Elodea	Water Lettuce
Eurasian Watermillfoil*	Water Hyacinth
Horned Pondweed*	Widgeon Grass
Hydrilla	Pondweed (e.g., Sago, American,)*

\* Variable control may be obtained in waters with greater calcium carbonate hardness.

### Timing of Treatments

When target vegetation is actively growing, apply Nautique Aquatic Herbicide to the area of greatest concentration of foliage in such a way as to evenly distribute the herbicide. In lakes, reservoirs, ponds, and static canals, the application site is defined by this label as the specific location where Nautique is applied. In slow moving and flowing canals and rivers, the application site is defined by this label as the target location for plant control. In order to maximize effectiveness, apply Nautique early in the day under bright or sunny conditions when water temperatures are at least 60 F (15 C). The activity of this product may be reduced if there is insufficient penetration of light into the water or if the plants and weeds are covered with silt, scale, or algae.

If algae mats are thick, use high pressure when spraying to break up the algae mats.

### Dissolved Oxygen Consideration

Treatment of aquatic plants and weeds can result in a reduction of dissolved oxygen due to the decomposition of the dead vegetation. This loss of dissolved oxygen can cause fish suffocation. To minimize this possible hazard treat 1/3 to 1/2 of the water area in a single operation, then wait 10-12 days before treating the remaining area. Begin treatment in the shallow areas, gradually proceeding outward in bands to permit the fish to move into the untreated area.

### Application Options

Nautique Aquatic Herbicide can be applied directly as a surface spray, subsurface through trailing weighted hoses, or in combination with other aquatic herbicides and algaecides, surfactants, sinking agents, polymers, or penetrants. These products are used to improve the retention time, sinking, and distribution of the herbicide. For surface application, this product may be applied diluted or undiluted, whichever is most suitable to insure uniform coverage of the area to be treated.

Aquatic plants and weeds will typically drop below the surface within 4-7 days after treatment. The complete results of treatment will be observed in 3-4 weeks in most cases. In heavily infested areas a second application may be necessary after 10-12 weeks. Repeating application of this product too soon after initial application may have no effect.

Use the lower rates for treating shallow water and the higher rates for treating deeper water and heavier infestations. Surface applications may be made from shore into shallow water along the shoreline.

Nautique Aquatic Herbicide inverts easily using either tank mix or multi-fluid mixer techniques. For submersed plants invert applications should be made through weighted hoses dragged below the water surface; for heavy infestations, direct application is preferable.

### No Restrictions on Water Use

Waters treated with Nautique may be used immediately after application for swimming, fishing, drinking, livestock watering, or irrigating turf and ornamental plants.

### Permits

Some states may require permits for the application of this product to public waters. Check with your local authorities.

### Application Rates

Recommended application rates in the chart below are based on minimal water flow in ponds, lakes, reservoirs, and irrigation conveyance or drainage systems. Treatments that extend chemical contact time with target vegetation will generally result in improved efficacy. In lakes, reservoirs, ponds, and static canals, the application

site is defined by this label as the specific location where Nautique is applied. In conveyance systems where significant water flow results in rapid off-site movement of copper, consult the Flowing Water Treatment Instructions for the recommended application instructions.

Application Rates		Gallons Per Surface Acre				Liters Per Surface hectare			
		Depth in Feet				Depth in meters			
Relative Density	ppm	1	2	3	4 <sup>1</sup>	0.5	0.75	1.0	1.25 <sup>2</sup>
Low Density	.5	1.5	3.0	4.5	6.0	12.0	24.1	36.1	48.2
	.6	1.8	3.6	5.4	7.2	14.9	29.8	44.7	59.6
Medium Density	.7	2.1	4.2	6.3	8.4	17.2	34.4	51.6	68.8
	.8	2.4	4.8	7.3	9.6	19.5	39.0	58.5	78.0
High Density	.9	2.7	5.4	8.1	10.8	21.8	43.6	65.4	87.2
	1.0 <sup>3</sup>	3.0	6.0	9.0	12.0	24.1	48.2	72.3	96.4

<sup>1</sup>For depths greater than 4 ft (1.25 m) add rates given for the sum of the corresponding depths in the chart

<sup>2</sup>Do not apply more than 1.0 ppm copper per application

**Free-Floating Plants** Apply Nautique at a rate of 8-12 gallons/acre for control of water hyacinth and salvinia and 4-6 gallons/acre for control of water lettuce. Add Nautique and appropriate surfactant to 100 gallons of water and use an adequate spray volume to insure good coverage of the plant.

## Tank Mix

### Nautique + Sonar A.S. Tank Mix (Except CA)

The following mixture can be used to provide rapid control of dense infestations of coontail, duckweed, egeria, elodea, Eurasian watermilfoil, hydrilla, sago and American pondweed, naiads, and other susceptible species. Apply 1 to 4 gallons of Nautique per surface acre in conjunction with normal Sonar rates. Observe all cautions and restrictions on the labels of both products used in this mixture.

### Nautique + Reward® Tank Mix

The following mixture can be used to enhance control of coontail, duckweed, egeria, elodea, Eurasian watermilfoil, hydrilla, pondweeds (Potamogeton species), salvinia, water lettuce, water hyacinth, and other susceptible species. Tank mix a ratio of 2:1 or 1.5:1 Nautique to Reward. This can be applied as a tank mix or metered in as a concentrate. The addition of a surfactant is recommended to enhance performance on floating plants. Observe all cautions and restrictions on the labels of both products used in this mixture. **DO NOT MIX CONCENTRATES IN TANK WITHOUT FIRST ADDING WATER.**

## Flowing Water Treatment :

### Drip System or Metering Pump Application for Canals, Ditches, and Laterals

This product should be applied as soon as submersed macrophytes begin to interfere with normal

delivery of water (clogging of lateral head gates, suction screens, weed screens, and siphon tubes). Delaying treatment could perpetuate the problem causing massing and compacting of plants. Heavy infestations and low flows may result in pooling or uneven chemical distribution resulting in unsatisfactory control. Under these conditions increasing the water flow rate during application may be necessary. In flowing canals the application site is defined by this label as the target location for aquatic plant control.

To achieve desired control with Nautique herbicide in flowing waters, it is recommended that a minimum exposure period of three hours be maintained. Other factors to consider include: plant species and density of infestation and water temperature and hardness. Treatment on bright sunny days will tend to enhance efficacy of this product.

1. Treatment with Nautique requires accurate calculations of water flow rates. Devices that provide accurate flow measurements such as weirs or orifices are the preferred method, however, the volume of water to be treated may also be estimated using the following formula:

$$\text{Average width (ft.)} \times \text{Average Depth (ft.)} \times \text{Average Velocity (ft./sec)} = \text{Cubic feet per Second (CFS)}$$

The velocity can be estimated by determining the length of time it takes a floating object to travel a defined distance. Divide the distance (ft.) by the time (sec.) to estimate velocity (ft./sec). This measure should be repeated 3 times at the intended application site and then calculate the average velocity.

2. After accurately determining the water flow rate in C.F.S. or gallons/minute, find the corresponding drip rate in the chart below.

Water Flow Rate		ppm Copper	Chemical Drip Rate	
C.F.S	Gal/Min.		Quart/ Hr	MI / min
1	450	0.5 – 1.0	0.5 – 1.0	8.0 – 16.0
2	900	0.5 – 1.0	1.0 – 2.0	16.0 – 32.0
3	1350	0.5 – 1.0	1.5 – 3.0	23.5 – 47.0
4	1800	0.5 – 1.0	2.0 – 4.0	31.5 – 63.0
5	2250	0.5 – 1.0	2.5 – 5.0	39.5 – 79.0

Calculate the amount of product needed to maintain the drip rate for a treatment period of 3 or more hours by multiplying quart/hr x 3; ml / min. by 180; or Fl. oz. / min x 180. Dosage will maintain 1.0 ppm copper concentration in the treated water for the treatment period. Introduction of the chemical should be made in the channel at weirs or other turbulence-creating structures to promote the dispersion of the chemical.

Pour the required amount of this product into a drum or tank equipped with a brass needle valve and constructed to maintain a constant drip rate. Use a stopwatch and appropriate measuring container to set the desired drip rate. Readjust accordingly if the canal flow rate changes during the treatment period. This product can also be applied by using metering pumps that adjust to flow rates in the canal.

Results can vary depending upon species and density of vegetation, desired distance of control and flow rate, and impact of water quality on copper residues and efficacy. Consult an Aquatic Specialist to determine optimal use rate and treatment period under local conditions. Periodic maintenance treatments may be required to maintain seasonal control.

### Irrigation Ponds

When applying to irrigation ponds, it is best to hold water for a minimum of 3 hours before irrigating to ensure proper exposure of Nautique at targeted rates to plants. If water is to be continually pumped from the treated system during application, application techniques (drip, injection, or multiple spray applications) should be made to compensate for dilution of Nautique within the targeted area.

### General Treatment Notes

The following suggestions apply to the use of this product as an algaecide or herbicide in all approved use sites. For optimum effectiveness:

- Apply early in the day under calm, sunny conditions when water temperatures are at least 60 deg. F.
- Treat when growth first begins to appear or create a nuisance, if possible.
- Apply in a manner that will ensure even distribution of the chemical within the treatment area.
- Re-treat areas if regrowth begins to appear and seasonal control is desired. Allow one to two weeks between consecutive treatments.
- Allow seven to ten days to observe the effects of treatment (bleaching and breaking apart of plant material).

### Storage and Disposal

Store in a cool, dry place.

**Pesticide Disposal:** Do not contaminate water, food or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incinerate, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## Warranty Disclaimer

SePRO Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. **SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

## Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to the label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation as the seller. All such risks shall be assumed by the buyer.

## Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies can not be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or Limitation of Remedies in any manner.